

# NUCLEAR SHAPE ISOMERS

P. MÖLLER and A. J. SIERK

Theoretical Division, Los Alamos National Laboratory  
Los Alamos, NM 87545

R. BENGTSSON

Department of Mathematical Physics, Lund Institute of Technology,  
P. O. Box 118, SE-22100 Lund, Sweden

H. SAGAWA

Center for Mathematical Sciences, University of Aizu  
Aizu-Wakamatsu, Fukushima 965-80, Japan

and

T. ICHIKAWA

Yukawa Institute for Theoretical Physics,  
Kyoto University, Kyoto 606-8502, Japan

Submitted to ATOMIC DATA AND NUCLEAR DATA TABLES May 10, 2010

Accepted September 14, 2010

We calculate potential-energy surfaces as functions of spheroidal ( $\epsilon_2$ ), hexadecapole ( $\epsilon_4$ ), and axial-asymmetry ( $\gamma$ ) shape coordinates for 7206 nuclei from  $A = 31$  to  $A = 290$ . We tabulate the deformations and energies of all minima deeper than 0.2 MeV *and* of the saddles between all pairs of minima. The tabulation is terminated at  $N = 160$ . Our study is based on the FRLDM macroscopic-microscopic model defined in ATOMIC DATA AND NUCLEAR DATA TABLES [59, 185 (1995)]. We also present potential-energy contour plots versus  $\epsilon_2$  and  $\gamma$  for 1224 even-even nuclei in the region studied. We can identify nuclei for which a *necessary* condition for *shape isomers* occurs, namely multiple minima in the calculated potential-energy surface. We find that the vast majority of nuclear shape isomers occur in the  $A = 80$  region, the  $A = 100$  region, and in a more extended region centered around  $^{208}\text{Pb}$ . A calculated region of shape isomers that has so far not been extensively explored is the region of neutron-deficient actinides “north-east” of  $^{208}\text{Pb}$ .